



Hexachloroethane: a highly efficient reagent for the synthesis of chlorosilanes from hydrosilanes

Veerachai Pongkittiphan^{a,b,c}, Emmanuel A. Theodorakis^c, Warinthorn Chavasiri^{a,b,*}

^a Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

^b Center for Petroleum, Petrochemicals, and Advance Materials, Chulalongkorn University, Bangkok 10330, Thailand

^c Department of Chemistry & Biochemistry, University of California, San Diego, La Jolla, CA 92093-0358, USA

ARTICLE INFO

Article history:

Received 15 March 2009

Revised 9 May 2009

Accepted 22 May 2009

Available online 27 May 2009

Keywords:

Hexachloroethane

Hydrosilanes

Chlorosilanes

Palladium(II) chloride

Chlorinating agents

ABSTRACT

A new and efficient chlorination protocol is presented for the preparation of chlorosilanes from hydrosilanes. A variety of chlorinating agents in combination with palladium(II) chloride as the catalyst are examined. Among them, hexachloroethane is found to be the best choice, furnishing the desired product in good to quantitative yields under mild conditions. Various hydrosilanes are used as starting materials to explore the scope of this reaction.

© 2009 Published by Elsevier Ltd.

Chlorosilanes play an important role in organic synthesis for the protection of highly reactive hydroxy and amino functional groups, as reagents for Mukaiyama aldol condensations,^{1–4} or as precursors of organochlorosilanes.^{5,6} In general, chlorosilanes are prepared by treatment of hydrosilanes with alkyl or allyl chlorides in the presence of CuCl₂,⁷ NiCl₂, PdCl₂^{8,9} or Pd/C¹⁰ as catalysts. However, these methods have certain limitations arising from the need for high temperatures, long reaction times or the use of expensive and toxic chlorinating agents.^{5,11} Recently, efficient systems were introduced for the conversion of alcohols into alkyl chlorides using the combination of PPh₃ and chlorinating agents such as Cl₃CCOCl₃, Cl₃CCN, Cl₃CCOOEt and Cl₃CCONH₂.^{12,13} Herein, we report a new and efficient procedure for the preparation of chlorosilanes from hydrosilanes using chlorinating agents in the presence of a catalytic amount of PdCl₂.

Several chlorinating agents were explored for the conversion of triisopropylhydrosilane (TIPS-H) into triisopropylsilyl chloride (TIPS-Cl) (Table 1). The yields of TIPS-Cl were quantified by ¹H NMR spectroscopy using toluene as an internal standard.

As shown in Table 1 (entry 1), PdCl₂ can transfer its chlorine atoms quantitatively to TIPS-H. However, the use of hexachloroacetone, trichloroacetonitrile and trichloroacetamide in combination with this catalyst produced TIPS-Cl in poor yields only (entries 3, 8 and 9, respectively). Of the three ethyl chloroacetates (entries 4, 6 and 7), Cl₂CHCOOEt provided the highest yield of TIPS-Cl, while

Cl₃CCOOEt, which has more chlorine atoms, gave a lower silyl chlorination yield. Comparison of the chloromethanes (entries 13–15) showed that CHCl₃ furnished a higher yield of chlorination product compared to CCl₄ or CH₂Cl₂. The product yield can be increased by carrying out these reactions at 55 °C for 3 h and 6 h, respectively (entries 3 and 4). The more reactive acetyl chloride afforded TIPS-Cl in quantitative yield (entry 2), however, due to its toxicity and acidity, it cannot be used in certain cases. On the other hand, comparison of different chloroethanes (entries 10–12) indicated that hexachloroethane afforded the highest yield of TIPS-Cl. In fact, 0.5 M equiv of this reagent led to quantitative conversion of TIPS-H into TIPS-Cl after only 1 h at 25 °C. When this reaction was repeated on a large scale, we were able to obtain a 99% yield of TIPS-Cl after vacuum distillation (98 °C, 20 mmHg).¹⁴

Table 2 reveals the effect of the amount and type of catalyst (PdCl₂ or Pd/C) on this reaction. As expected, no reaction occurred when the Pd catalyst was not present demonstrating that the catalyst is essential (entry 1). Moreover, the product yield was reduced to 81% when 0.5 mol % PdCl₂ was employed (entry 3).


The results of the optimization studies on the amount of Cl₃CCCl₃ required for this chlorination procedure are shown in Table 3. Use of 0.17 M equiv of this reagent (corresponding to 1 equiv of chlorine atoms) under standard reaction conditions produced only a 69% yield of TIPS-Cl (entry 2). The yield of TIPS-Cl was increased to 95% by using Cl₃CCCl₃ (0.30 mmol) (1.8 equiv of chlorine atoms). The use of 0.5 mmol of hexachloroethane (3 equiv of chlorine atoms) resulted in a quantitative yield of TIPS-Cl after only 30 min at room temperature (entry 5).

* Corresponding author. Tel.: +66 2 218 7625; fax: +66 2 218 7598.

E-mail address: warintho@yahoo.com (W. Chavasiri).

Table 1


The effect of chlorinating agents on the conversion of TIPS-H into TIPS-Cl




Entry	Chlorinating agent	Equivalents (mmol)	% Yield ^a Si-Cl	% Recovery Si-H
1	None	—	2	98
2	CH ₃ COCl	0.75	69	31
		1.50	Quant	0
3	Cl ₃ CCOCl ₃	0.50	17 (59) ^b (92) ^c	83 (41) ^b (8) ^c
4	Cl ₃ CCOOEt	0.50	15 (39) ^b (35) ^c	85 (61) ^b (65) ^c
5	Cl ₃ CCOOH	0.50	18	82
6	Cl ₂ CHCOOEt	0.50	83	17
7	ClCH ₂ COOEt	0.50	10	90
		1.50	24	76
8	Cl ₃ CCN	0.50	11	89
9 ^d	Cl ₃ CCONH ₂	0.50	15	85
10	ClCH ₂ CH ₂ Cl	0.50	16	84
11	Cl ₂ CHCHCl ₂	0.50	32	68
12	Cl ₃ CCCl ₃	0.50	Quant (99) ^{e,f}	0
13	CCl ₄	0.50	18	82
14	CHCl ₃	0.50	65	35
15	CH ₂ Cl ₂	0.50	14	86
16	Cl ₃ CCH ₃	0.50	54	46

^a % yield was determined by ¹H NMR using toluene as an internal standard.^b The reaction was carried out at 55 °C for 3 h.^c The reaction was carried out at 55 °C for 6 h.^d Not completely soluble.^e 20 mmol scale reaction was performed with 0.5 mol % PdCl₂.^f Isolated by distillation (bp 98 °C, 20 mmHg).**Table 2**

The effect of the amount and type of catalyst on the conversion of TIPS-H into TIPS-Cl




Entry	Catalyst (% mmol)	% Yield Si-Cl	% Recovery Si-H
1	None	—	100
2	PdCl ₂ (1.0%)	Quant	0
3	PdCl ₂ (0.5%)	81	19
4	Pd/C (1.0%)	Quant	0

Table 3The effect of the amount of Cl₃CCCl₃ on the conversion of TIPS-H into TIPS-Cl

Entry	Cl ₃ CCCl ₃ (mmol)	% Yield Si-Cl	% Recovery Si-H
1	0.125	48	52
2	0.17	69	31
3	0.25	90	10
4	0.30	93	7
5 ^a	0.50	Quant	0
		Quant	0
6	0.75	Quant	0

^a Reaction time: 30 min.

A general mechanism can be proposed (Scheme 1) which proceeds by the interaction of the chlorinating agent (RCl) with Pd(0) by oxidative addition, followed by transmetalation and reductive elimination to regenerate Pd(0).¹⁵


**Scheme 1.** A general mechanistic pathway for the conversion of hydrosilanes into chlorosilanes.

This method is applicable to the synthesis of various chlorosilanes (Table 4). In the case of liquid hydrosilanes the chlorination proceeded best under neat conditions, but with solid hydrosilanes, for example, Ph₃SiH, a solvent was required. Among the various solvents examined, THF was found to produce consistently higher chlorination yields. As shown in Table 4, this reaction can be applied to various hydrosilanes including monochloro diphenyl hydrosilane (entries 15–18). The reaction was performed under the optimized conditions: Cl₃CCCl₃ (0.25 mmol) and 1% PdCl₂ at room temperature for one hour without any solvent unless otherwise stated. All the selected hydrosilanes were transformed successfully into the corresponding chlorosilanes with experimental modifications in some cases.

In conclusion, hexachloroethane was found to be an efficient chlorinating agent for the conversion of hydrosilanes into chlorosilanes in the presence of PdCl₂ as the catalyst. This chlorination

Table 4

The conversion of monohydrosilanes into chlorosilanes



Entry	Hydrosilane	Solvent (0.25 mL)	% Yield Si-Cl	% Recovery Si-H
1	Et ₃ SiH	—	78	22
2 ^a		—	Quant (99.8) ^{fg}	0
3	(Me ₃ Si) ₃ SiH	—	Quant	0
4		THF	Quant	0
5	Ph ₃ SiH	THF	Quant	0
6	Ph(Me) ₂ SiH	THF	30	70
7 ^a		THF	66	34
8 ^{a,c}		THF	55	45
9 ^d		THF	61	39
10 ^e		THF	60	40
11 ^b	<i>t</i> -Bu(Me) ₂ SiH	—	67	33
12 ^a		THF	86	14
13 ^{a,c}		THF	89	11
14		THF	80	20
15	Ph ₂ SiClH	—	89	11
16		THF	77	23
17 ^a		THF	83	17
18 ^{a,c}		THF	78	22

^a Using 0.50 mmol of Cl₃CCCl₃.^b The reaction was carried out in an ice bath.^c The reaction was run for 2 h.^d Using 0.75 mmol of Cl₃CCCl₃.^e Using 1.00 mmol of Cl₃CCCl₃.^f 20 mmol scale reaction was performed with 0.5 mol % PdCl₂.^g Isolated by vacuum distillation (bp 45–47 °C, 20 mmHg).

reaction proceeds cleanly and rapidly under mild conditions and, in certain cases, in the absence of any solvent. Sub-equivalent amounts of hexachloroethane are needed for quantitative chlorination (typically between 0.25 and 0.5 equiv), which together with the fact that this compound is commercially available and inexpensive, renders this reaction attractive as a general method for the synthesis of chlorosilanes.

Acknowledgements

We gratefully thank the Natural Products Research Unit, Department of Chemistry, Faculty of Science, Chulalongkorn University, for the provision of chemicals and laboratory facilities. Financial support from TRF Master Research Grants, the Graduate school, Chulalongkorn University and the National Institutes of Health (Grant GM081484) is also acknowledged.

References and notes

1. Birkofer, L.; Stuhl, O.; Patai, S.; Rappoport, Z. *The Chemistry of Organic Silicon Compounds*; Wiley: Chichester, 1989. Chapter 10, pp. 655–761.
2. Larson, G. L.; Patai, S.; Rappoport, Z. *The Chemistry of Organic Silicon Compounds*; Wiley: Chichester, 1989. Chapter 11, pp. 763–888.
3. Cunico, R. F.; Bedell, L. J. *Org. Chem.* **1980**, *45*, 4797–4798.
4. Khalafi-Nezhad, A.; Alamdari, R. F.; Zekri, N. *Tetrahedron* **2000**, *56*, 7503–7506.
5. Masaoka, S.; Banno, T.; Ishikawa, M. *J. Organomet. Chem.* **2006**, *691*, 174–181.
6. Kobayashi, T.; Pannell, K. H. *Organometallics* **1991**, *10*, 1960–1964.
7. Kunai, A.; Sakurai, T.; Toyoda, E.; Ishikawa, M. *J. Organomet. Chem.* **1996**, *15*, 2478–2482.
8. Kunai, A.; Ohshita, J. *Organometallic* **2003**, *22*, 686, 3–15.
9. (a) Ferreri, C.; Costantino, C.; Chatgililoglu, C.; Boukherroub, R.; Manuel, G. J. *Organomet. Chem.* **1998**, *554*, 135–137; (b) Ferreri, C.; Costantino, C.; Romeo, R.; Chatgililoglu, C. *Tetrahedron Lett.* **1999**, *40*, 1197–1200.
10. (a) Sommer, L. H.; Citron, J. D. *J. Org. Chem.* **1967**, *32*, 2470–2472; (b) Citron, J. D.; Lyons, J. E.; Sommer, L. H. *J. Org. Chem.* **1968**, *33*, 638–640.
11. Citron, J. D. *J. Org. Chem.* **1969**, *34*, 1977–1979.
12. Pluempanupat, W.; Chavasiri, W. *Tetrahedron Lett.* **2006**, *47*, 6821–6823.
13. (a) Magid, R. M.; Fruchey, O. S.; Johnson, W. L.; Allen, T. G. *J. Org. Chem.* **1979**, *44*, 359–363; (b) Jang, D. O.; Park, D. J.; Kim, J. *Tetrahedron Lett.* **1999**, *40*, 5323–5326.
14. *Typical experimental for the conversion of hydrosilanes into chlorosilanes*: To a stirred mixture of hexachloroethane (2.37 g, 10 mmol, 0.5 equiv) and Pd(II)Cl₂ (18.5 mg, 0.1 mmol, 0.5 mol %) at room temperature under an N₂ atmosphere was added triisopropylsilane (3.16 g, 20 mmol, 1 equiv) over a period of 2 min. *Note*: the reaction is exothermic and large-scale reactions should be cooled using an ice bath. The reaction mixture became homogeneous after 5 min and was allowed to stir for 1 h. The crude residue was then distilled under vacuum (bp 98 °C, 20 mmHg) to yield triisopropylsilyl chloride (3.78 g, 19.7 mmol, 99%). In a similar manner triethylsilane (2.33 g, 20 mmol) was converted into triethylsilyl chloride (3.0 g, 19.96 mmol, quant. after distillation at 45–47 °C, 20 mmHg).
15. Cho, Y. S.; Han, J. S.; Yoo, B. R.; Kang, S. O.; Jung, I. N. *Organometallics* **1998**, *17*, 570–573.